

The Blow-up Lemma and growing degrees

Andreas Würfl

Technische Universität München

Combinatorics Conference in Lisbon

(joint work with Julia Böttcher, Yoshi Kohayakawa & Anusch Taraz)

- 1 The Blow-up Lemma

- 2 Growing degrees
 - Arrangeability of graphs
 - Random greedy embedding

- 3 Making things rigorous
 - Auxiliary graphs
 - Matchings in auxiliary graphs

Let $\varepsilon, \delta > 0$. The graph $(A \dot{\cup} B, E)$ with $|A| = |B| = n$ is an (ε, δ) -super-regular pair if


- $|d(X, Y) - d(A, B)| \leq \varepsilon$ for all $X \subseteq A, Y \subseteq B$ with $|X|, |Y| \geq \varepsilon n$,
- $\deg(v) \geq \delta n$ for all $v \in A \cup B$.

Definition

(ε, δ) -SUPER-REGULAR

Let $\varepsilon, \delta > 0$. The graph $(A \dot{\cup} B, E)$ with $|A| = |B| = n$ is an (ε, δ) -super-regular pair if

- $|d(X, Y) - d(A, B)| \leq \varepsilon$ for all $X \subseteq A, Y \subseteq B$ with $|X|, |Y| \geq \varepsilon n$,
- $\deg(v) \geq \delta n$ for all $v \in A \cup B$.



- regularity \leftrightarrow
“densities as expected”
- high minimum degree

For $\delta > 0$, $\Delta \in \mathbb{N}$ there is $\varepsilon > 0$ such that the following holds. If H has $\Delta(H) \leq \Delta$ and $H \subseteq K_{n,n}$ then H is subgraph of any (ε, δ) -super-regular pair $G = (A \dot{\cup} B, E)$ with $|A| = |B| = n$.


another proof was given by:

RÖDL, RUCIŃSKI, '99

For $\delta > 0$, $\Delta \in \mathbb{N}$ there is $\varepsilon > 0$ such that the following holds. If H has $\Delta(H) \leq \Delta$ and $H \subseteq K_{n,n}$ then H is subgraph of any (ε, δ) -super-regular pair $G = (A \dot{\cup} B, E)$ with $|A| = |B| = n$.

another proof was given by:


RÖDL, RUCIŃSKI, '99



For $\delta > 0$, $\Delta \in \mathbb{N}$ there is $\varepsilon > 0$ such that the following holds. If H has $\Delta(H) \leq \Delta$ and $H \subseteq K_{n,n}$ then H is subgraph of any (ε, δ) -super-regular pair $G = (A \dot{\cup} B, E)$ with $|A| = |B| = n$.

another proof was given by:


RÖDL, RUCIŃSKI, '99



For $\delta > 0$, $\Delta \in \mathbb{N}$ there is $\varepsilon > 0$ such that the following holds. If H has $\Delta(H) \leq \Delta$ and $H \subseteq K_{n,n}$ then H is subgraph of any (ε, δ) -super-regular pair $G = (A \dot{\cup} B, E)$ with $|A| = |B| = n$.

another proof was given by:

RÖDL, RUCIŃSKI, '99



The Blow-up Lemma for growing degrees

Theorem

KOMLÓS, SÁRKÖZY, SZEMERÉDI, '97

For $\delta > 0$, $\Delta \in \mathbb{N}$ there is $\varepsilon > 0$ such that the following holds. If H has $\Delta(H) \leq \Delta$ and $H \subseteq K_{n,n}$ then H is subgraph of any (ε, δ) -super-regular pair $(A \dot{\cup} B, E)$ with $|A| = |B| = n$.

The Blow-up Lemma for growing degrees

Theorem

BÖTTCHER, KOHAYAKAWA, TARAZ, W. '11

For $\delta > 0$, $a \in \mathbb{N}$ there is $\varepsilon > 0$ such that the following holds. If H has $\Delta(H) \leq \sqrt{n}/\log n$ and $H \subseteq K_{n,n}$ is a -arrangeable then H is subgraph of any (ε, δ) -super-regular pair $(A \dot{\cup} B, E)$ with $|A| = |B| = n$.


The Blow-up Lemma for growing degrees

Theorem


BÖTTCHER, KOHAYAKAWA, TARAZ, W. '11

For $\delta > 0$, $a \in \mathbb{N}$ there is $\varepsilon > 0$ such that the following holds. If H has $\Delta(H) \leq \sqrt{n}/\log n$ and $H \subseteq K_{n,n}$ is a -arrangeable then H is subgraph of any (ε, δ) -super-regular pair $(A \dot{\cup} B, E)$ with $|A| = |B| = n$.

$G = (V, E)$ is a -arrangeable if there is an ordering $V = \{v_1, \dots, v_n\}$ with $|N_L(N_R(v_i))| \leq a$ for all $i = 1, \dots, n$.



Examples of arrangeable graphs




- trees are 1-arrangeable
- planar graphs are 761-arrangeable
- planar graphs are 10-arrangeable
- graphs without a K_p -subdivision are p^8 -arrangeable

CHEN, SHELP '93

KIERSTEAD, TROTTER '94


RÖDL, THOMAS '94

Sketch of proof: a random greedy embedding




(ε, δ) -super-regular

Sketch of proof: a random greedy embedding




Sketch of proof: a random greedy embedding




$$C(x) = \bigcap_{y \in N_L(x)} N_G(\varphi(y))$$

Sketch of proof: a random greedy embedding




$$C(x) = \bigcap_{y \in N_L(x)} N_G(\varphi(y))$$

Sketch of proof: a random greedy embedding




$$C(x) = \bigcap_{y \in N_L(x)} N_G(\varphi(y))$$

Sketch of proof: a random greedy embedding




$$C(x) = \bigcap_{y \in N_L(x)} N_G(\varphi(y))$$

Sketch of proof: a random greedy embedding




$$C(x) = \bigcap_{y \in N_L(x)} N_G(\varphi(y))$$

Sketch of proof: a random greedy embedding




MESSAGE: respect your successors!


Sketch of proof: a random greedy embedding




Sketch of proof: a random greedy embedding



Sketch of proof: a random greedy embedding




Sketch of proof: a random greedy embedding



Does this always work?




Respect one successor...

...with the help of ε -regularity



Respect one successor...


...with the help of ε -regularity



(ε, δ) -super-regular


Respect one successor...

...with the help of ε -regularity



Respect one successor...

...with the help of ε -regularity



(ε, δ) -super-regular

all but $2\varepsilon n$ vertices in $C(3)$ have “correct” degree into $C(4)$

Respect all successors. . .

. . . even if their number is growing with n

Respect all successors. . .

. . . even if their number is growing with n


Problem: each successor might “kill” $\varepsilon' n$ candidates

Respect all successors. . .

. . . even if their number is growing with n

Problem: each successor might “kill” $\varepsilon' n$ candidates

Solution: the *a-arrangeability* of H




Respect all successors. . .

. . . even if their number is growing with n

Problem: each successor might “kill” $\varepsilon' n$ candidates

Solution: the a -arrangeability of H




all successors of v_i have at most a predecessors in total


\Rightarrow we have to respect at most 2^a different candidate sets

\Rightarrow we loose at most $2^a \varepsilon' n$ candidates

The auxiliary graphs



The auxiliary graphs



$$F_1 = (X_1 \cup V_1, E_1) \text{ with } \{x, v\} \in E_1 \text{ iff } v \in C(x)$$

One auxiliary graph, two nice properties

Lemma

With probability $1/2$ all auxiliary graphs F_i inherit *some* regularity from G .
 \Rightarrow random embedding algorithm gives an **almost spanning embedding**.

One auxiliary graph, two nice properties

Lemma

With probability $1/2$ all auxiliary graphs F_i inherit *some* regularity from G .
 \Rightarrow random embedding algorithm gives an **almost spanning embedding**.

One auxiliary graph, two nice properties

Lemma

With probability $1/2$ all auxiliary graphs F_i inherit *some* regularity from G .
 \Rightarrow random embedding algorithm gives an **almost spanning embedding**.

- preprocessing:
select an arrangeable ordering with a stable set at its end

One auxiliary graph, two nice properties

Lemma

With probability $1/2$ all auxiliary graphs F_i inherit *some* regularity from G .
 \Rightarrow random embedding algorithm gives an **almost spanning embedding**.

- preprocessing:
select an arrangeable ordering with a stable set at its end

Lemma

All auxiliary graphs have high min. degree with probability tending to 1.

One auxiliary graph, two nice properties

Lemma

With probability $1/2$ all auxiliary graphs F_i inherit *some* regularity from G .
 \Rightarrow random embedding algorithm gives an **almost spanning embedding**.

- preprocessing:
select an arrangeable ordering with a stable set at its end

Lemma

All auxiliary graphs have high min. degree with probability tending to 1.

- \exists **perfect matching** in the “remaining part” of each auxiliary graph

One auxiliary graph, two nice properties

Lemma

With probability $1/2$ all auxiliary graphs F_i inherit *some* regularity from G .
 \Rightarrow random embedding algorithm gives an **almost spanning embedding**.

- preprocessing:
select an arrangeable ordering with a stable set at its end

Lemma

All auxiliary graphs have high min. degree with probability tending to 1.

- \exists **perfect matching** in the “remaining part” of each auxiliary graph
- these perfect matchings define a **spanning embedding**.

The 'obvious' next step

$G = (V, E)$ is d -degenerate if there is an ordering $V = \{v_1, \dots, v_n\}$ with $|N_L(v_i)| \leq d$ for all $i = 1, \dots, n$.



The 'obvious' next step

$G = (V, E)$ is d -degenerate if there is an ordering $V = \{v_1, \dots, v_n\}$ with $|N_L(v_i)| \leq d$ for all $i = 1, \dots, n$.



Replace *const. max. degree* by d -degenerate in the Blow-up Lemma!

The 'obvious' next step

$G = (V, E)$ is d -degenerate if there is an ordering $V = \{v_1, \dots, v_n\}$ with $|N_L(v_i)| \leq d$ for all $i = 1, \dots, n$.



Replace *const. max. degree* by d -degenerate in the Blow-up Lemma!

Far goal

For $\delta, \mu > 0$, $d \in \mathbb{N}$ there is $\varepsilon, \xi > 0$ such that the following holds. If H has $\Delta(H) \leq \xi n$ and $H \subseteq K_{(1-\mu)n, (1-\mu)n}$ is d -degenerate then H is subgraph of any (ε, δ) -super-regular pair $(A \dot{\cup} B, E)$ with $|A| = |B| = n$.