



Stochastic Programming

Prof. Dr. Raymond Hemmecke | Silvia Lindner

Exercise sheet 4

Exercise 4.1, continued Benders' decomposition: Consider the **facility location problem**: A company wants to expand into new territories and considers m possible locations for new factories, the cost of opening a factory at location i is f_i . A market analysis has revealed the demand for each of the n potential customers, the cost of supplying customer j from factory i is c_{ij} . We want to determine at which locations a factory should be opened and which fraction of each customer's demand should be supplied from which factory in order to minimize the total cost (delivery plus factory setup).

Use variables $x_{ij} \in [0; 1]$ to determine which fraction of customer j 's demand should be supplied from the factory at location i and variables $y_i \in \{0, 1\}$ to determine whether or not to open a factory at location i .

1. Design an IP model for the facility location using the notation given above.
2. We want to apply Benders' decomposition to the problem. Formulate the dual subproblem for fixed y -variables. How can it be solved? Outline how to find a feasibility cut and an optimality cut.

Exercise 4.2 The L-Shaped Method: Consider Step 3 of Iteration 1 within Example 1.

a) Let $\xi = \xi_1$, $x = x^1$ and

$$\begin{aligned} w &= \min -24y_1 - 28y_2 \\ 6y_1 + 10y_2 &\leq 2400 \\ 8y_1 + 5y_2 &\leq 1600 \\ 0 &\leq y_1 \leq 500 \\ 0 &\leq y_2 \leq 100. \end{aligned}$$

i) Check that the optimal dictionary (the simplex tableau in the optimal solution) is

$$\begin{aligned} w &= -6100 + 3s_2 + 13s_4 \\ s_1 &= 575 + 6/8s_2 + 50/8s_4 \\ y_1 &= 137.5 - 1/8s_2 + 5/8s_4 \\ s_3 &= 362.5 + 1/8s_2 - 5/8s_4 \\ y_2 &= 100 - s_4 \end{aligned}$$

for slack variables s_1, s_2, s_3, s_4 .

ii) Check that this dictionary corresponds to the solution stated in Example 1.

- iii) Check that the optimal value $w = -6100$ is also attained by the dual variables.
- b) For $\xi = \xi_1$, the optimal solution is $w_1 = -6100$ and for $\xi = \xi_2$, the optimal solution is $w_2 = -8384$.
- i) Check that $w^1 = 0.4w_1 + 0.6w_2$.
- ii) Prove by linear programming duality that $w = \sum_{k=1}^K p_k w_k$, where w_k denotes the solution of the second-stage program, for realization k of ξ , $k = 1, \dots, K$.

Exercise 4.3 Consider the following problem:

$$\begin{aligned}
 w = \min \quad & 7x_1 + 11x_2 + E_\xi(q_1y_1 + q_2y_2) \\
 & y_1 + 2y_2 \geq d_1 - x_1 \\
 & y_1 \geq d_2 - x_2 \\
 & 0 \leq x_1 \leq 10 \\
 & 0 \leq x_2 \leq 10 \\
 & y_1, y_2 \geq 0.
 \end{aligned}$$

where $\xi^\top = (q_1, q_2, d_1, d_2)$ takes on the values $(26, 16, 6, 12)$ and $(14, 24, 10, 4)$ with probability 0.5 each.

In this example, the L-Shaped method selects $x^1 = (0, 0)^\top$ as starting point. The L-Shaped method can however be used with any other reasonable starting point. Take $x^1 = (1, 5)^\top$ as starting point and show that exactly the same steps are taken if the starting point is any point within the region $4 \leq x_2 \leq 6 + x_1$.